

V440 Per: the longest period overtone Cepheid

R. Baranowski^{1*}, R. Smolec², W. Dimitrov¹, T. Kwiatkowski¹,
A. Schwarzenberg-Czerny^{1,2†}, P. Bartczak¹, M. Fagas¹, W. Borczyk¹,
K. Kamiński¹, P. Moskalik², R. Ratajczak¹ and A. Rożek¹

¹ *Astronomical Observatory of Adam Mickiewicz University, ul. Słoneczna 36, PL 60-286 Poznań, Poland,*

² *Copernicus Astronomical Centre, ul. Bartycka 18, PL 00-716 Warsaw, Poland*

Accepted Received; in original form

ABSTRACT

V440 Per is a Population I Cepheid with the period of 7.57 day and low amplitude, almost sinusoidal light and radial velocity curves. With no reliable data on the 1st harmonic, its pulsation mode identification remained controversial. We obtained a radial velocity curve of V440 Per with our new high precision and high throughput Poznań Spectroscopic Telescope. Our data reach the accuracy of 130 m/s per individual measurement and yield a secure detection of the 1st harmonic with the amplitude of $A_2 = 140 \pm 15$ m/s. The velocity Fourier phase ϕ_{21} of V440 Per is inconsistent at the 7.25 σ level with those of the fundamental mode Cepheids, implying that the star must be an overtone Cepheid, as originally proposed by Kienzie et al. (1999). Thus, V440 Per becomes the longest period Cepheid with the securely established overtone pulsations. We show, that the convective nonlinear pulsation hydrocode can reproduce the Fourier parameters of V440 Per very well. Requirement to match the observed properties of V440 Per constrains free parameters of the dynamical convection model used in the pulsation calculations, in particular the radiative losses parameter.

Key words: stars: variables: Cepheids – stars: oscillations – stars: individual: V440 Per – techniques: spectroscopic – methods: data analysis – hydrodynamics

1 INTRODUCTION

The Population I *sinusoidal* or s-Cepheids are a small group of the Cepheids pulsating in the first radial overtone. In the Galaxy, where individual Cepheid distances are usually not accurately known, the s-Cepheids are discriminated from the fundamental mode pulsators with the Fourier decomposition of their light curves (Antonello et al. 1990). The method works well only for the variables with periods below 5 days. Fortunately, it can be extended to longer periods with the help of the radial velocity curves, as was shown by Kienzie et al. (1999). Studies of the longer period overtone Cepheids are of the great interest. The velocity Fourier parameters of the s-Cepheids display a very characteristic progression with period, which is attributed to the 2:1 resonance at $P = 4.2 - 4.6$ day between the first and fourth overtones (Kienzie et al. 1999; Feuchtinger et al. 2000). Unfortunately, any detailed modeling of this progression and pinpointing of the resonance position is hampered by a scarcity of the s-Cepheids with $P > 5.5$ day, with MY Pup being the only secure identification. One more can-

didate, V440 Per ($P = 7.57$ day), has been identified by Kienzie et al. (1999). They noted, that the velocity Fourier phase ϕ_{21} places this Cepheid away from the fundamental mode sequence and possibly onto the first overtone sequence. On this basis, Kienzie et al. (1999) proposed that V440 Per is an overtone pulsator. This hypothesis was further supported by determination of the phase-lag between the light curve and the radial velocity curve, $\Delta\Phi_1 = \phi_1^{Vr} - \phi_1^{mag}$, which placed V440 Per away from the fundamental sequence as well (Ogloza et al. 2000). However, V440 Per is a small amplitude, nearly sinusoidal variable. Even the best then available radial velocity data (Burki & Benz 1982) yielded large errors of the 1st harmonics Fourier parameters. The measurement error of the phase-lag $\Delta\Phi_1$ was large, too. This, and inconsistency with their hydrodynamic pulsation models led Szabó et al. (2007) to dispute the mode identification of V440 Per. They noted that the membership of this Cepheid in the fundamental mode sequence could not be rejected at the 3 σ confidence level. They argued that V440 Per is not an overtone pulsator, but rather a fundamental mode Cepheid of a very low amplitude.

In the present paper we report results of an extensive campaign of the observations of V440 Per with the Poznań Spectroscopic Telescope, lasting almost a year and aiming

* Based on observations from Poznań Spectroscopic Telescope

† e-mail alex@camk.edu.pl

Table 1. PST radial velocities of V440 Per. Full table in the electronic form only.

MJD	$\Delta v_r^{a)}$	$\Delta v_{r,corr}^{b)}$
54327.0490	-0.61	-0.52
54332.0517	3.04	3.14
54332.0683	2.96	3.06
...		
54649.0009	2.28	2.12
54650.9302	2.75	2.57
54650.9419	2.76	2.58

a) zero shifted by about -26.3 ± 0.2 km/s

b) corrected for instrumental effects (see text)

at obtaining high quality data suitable for the detailed diagnostic and comparison with the nonlinear models. In Sect. 2 we describe our instrument and observations. In Sect. 3 we discuss quality of our radial velocities and derive Fourier parameters of V440 Per. Comparison with Fourier parameters of other Galactic Cepheids and identification of the pulsation mode of V440 Per is discussed in Sect. 4. In Sect. 5 we perform detailed comparison of V440 Per with the nonlinear overtone Cepheid models. Our conclusions are summarized in Sect. 6.



2 OBSERVATIONS AND DATA REDUCTION

Our observations were obtained with the new Poznań Spectroscopic Telescope (PST) of Adam Mickiewicz University. Its full description is not published yet, hence we devote some space here to instrument description and data quality evaluation.

2.1 Poznań Spectroscopic Telescope

PST is located in Poland at Borowiec station, 20 km south from Poznań city, at a meagre elevation of 123 m above sea level. The PST consists of parallel twin 0.4 m Newton telescopes of the f-ratio 4.5, fixed on a single parallactic fork mount. An acquisition box at Newton focus of each telescope holds the tip of the fiber feeding our spectrograph, the Thor/Argon calibration lamp and the autoguider camera SBIG ST-7.

The telescope feeds via a fiber an Echelle spectrograph, a clone of MUSICOS design (Baudrand & Bohm 1992), red arm only. Our spectra are recorded with the low noise Andor DZ436 camera fitted with $2k \times 2k$ E2V 42-40 back illuminated CCD chip, cooled with Peltier cells. About 60 orders are recorded, covering a spectral range of 4480–9250 Å at the inverse resolution of $\lambda/\delta\lambda = 35000$. The spectrograph is located in a thermo-isolated enclosure in the telescope dome. The sliding-roof dome, the telescope and its spectrograph all operate under full computer control. A full description of our instrumentation will be published elsewhere. Our system is operated interactively from the terminal. Such an operation mode does not compromise our data quality but requires excessive work-load. A substantial software effort is needed to achieve the fully robotic operation.

**Figure 1.** Radial velocities of V440 Per *vs.* phase. First and second order Fourier fits are plotted with dashed and solid lines, respectively. The error bar indicates ± 1 standard deviation of residuals. The velocities are corrected for small instrumental effects (see text). Zero point is arbitrary.**Figure 2.** Top: residuals of the first order Fourier fit of V440 Per radial velocities (dashed line in Fig. 1) *vs.* phase. 1st harmonic is very small ($A_2 = 140 \pm 15$ m/s), yet clearly visible. Bottom: residuals of the second order Fourier fit *vs.* time.

2.2 Data pipeline

Routine CCD reductions up to spectra extraction, wavelength calibration and velocity measurement are performed using IRAF tasks combined into our reduction pipeline. Velocities are measured by cross-correlation, using IRAF FXCOR task. The internal error estimates from FXCOR, of order of few km/s, relate to the line width and not to the actual measurement precision. They may serve for weighting purposes, though. So far, we employed no standard star calibrations and our velocities are measured solely with respect to the Th/Ar lamp. For short span observations such a primitive procedure still yields root-mean-square (RMS) residuals in the 100–200 m/s range, for stars brighter than 11 mag. As demonstrated in Sect. 3, for intensively observed,

Table 2. Fourier parameters of V440 Per radial velocity curve

Name	Value	Error	Unit
T_0	54498.783	0.008	MJD
P	7.5721	0.0006	d
A_1	2.480	0.015	km/s
A_2	0.140	0.015	km/s
R_{21}	0.056	0.006	1
ϕ_{21}	2.759	0.117	rad

strictly periodic stars, any long-term effects of a floating instrumental zero-point may be removed with some assurance, yielding again RMS residuals less than 150 m/s, over one year span of data.

2.3 Observations

In total 158 radial velocity measurements of V440 Per were obtained with PST from 2007 August 15 till 2008 July 03. To reach signal-to-noise ratio of 70 we exposed spectra for 10–15 min and up to several spectra were obtained per night. The observed velocities are listed in Table 1 published in the electronic form. Phase-folded data with fitted Fourier series (Sect. 3) are displayed in Fig. 1. In Fig. 2 we plot residuals of the first order and the second order fits *vs.* pulsation phase and *vs.* time. Inspection of the plots demonstrates that our coverage was reasonably uniform, both in time and in frequency. From the residuals we estimate standard error of our individual measurements as 130 m/s.

3 FOURIER PARAMETERS

The plot of the phase-folded radial velocities reveals near sinusoidal variations with peak-to-peak amplitude of 5 km/s. Some data consistency checks are due prior to drawing any final conclusions. Cepheid phases and frequencies are known to vary slightly. Additionally, the instrument stability over 9 months needs checking, too. A preliminary nonlinear least squares fit of our data with the Fourier series of three harmonics terms (third order fit) yielded no significant 2nd harmonics of the main frequency nor the period derivative term. Our fitted frequency of 0.13206 ± 0.00001 c/d is less accurate, yet consistent within errors with the frequency derived by combining our data with the earlier measurements of Burki & Benz (1982), Arellano Ferro (1984) and Gorynya et al. (1992, 1996, 1998). Comparison of the data shows, that the weighted zero point shift of our velocities with respect to the previous authors is -26.3 ± 0.2 km/s.

The RMS deviation from the Fourier fit of all our data was 164 m/s, well in excess of 135 m/s obtained for the first part of the dataset. Worse, inspection of the residuals plotted against time sometimes revealed non-gaussian, bi-modal distribution. The origin of both effects seems to be instrumental. To confirm that, we expanded our Fourier model by including two linear terms proportional to the time interval from the mid-epoch and to the hour angle of the star at the moment of observation. The fitted values of these instrumental correction coefficients were significant at 6σ and 5σ level, respectively. From the overall covariance matrix we find maximum absolute value of the correlation coefficients




Figure 3. Fourier phase ϕ_{21} *vs.* pulsation period for Cepheid radial velocity curves. Fundamental mode Cepheids displayed with asterisks, except for low amplitude ones ($A_1 < 10$ km/s) marked with open triangles. Overtone Cepheids displayed with filled circles. Observational data points taken from Kienzie et al. (1999) and Moskalik et al. (2009). Straight line indicates the best fit to the segment of the fundamental mode progression. V440 Per displayed with open circle.

of 0.30, consistent with little interference between different fitted terms. These instrumental corrections reached up to ± 100 m/s. At this stage, we have no explanation for these corrections.

Our final second order Fourier fit, supplemented with the instrumental correction terms, yielded RMS deviation of 130 m/s, consistent with that obtained from the short-span observations. The values of the Fourier parameters of V440 Per radial velocity curve are listed in Table 2 (for exact formulae defining Fourier parameters and their errors see Appendix A).

4 PULSATION MODE OF V440 PER

Pulsation mode of a Cepheid can be established by measuring Fourier phase ϕ_{21} of its light curve (Antonello et al. 1990) or radial velocity curve (Kienzie et al. 1999). This Fourier parameter does not depend on the pulsation amplitude of the star and for each mode it follows a different, *tightly defined* progression with the pulsation period. For periods at which the two ϕ_{21} progressions are well separated, secure mode identification can be achieved.

In Fig. 3 we plot velocity ϕ_{21} of short period Galactic Cepheids against their pulsation period P . Fundamental mode pulsators and overtone pulsators are displayed with different symbols. V440 Per is plotted with an open circle. It is immediately obvious, that it is located far apart from the fundamental mode progression. This notion can be put on a quantitative basis. We selected a sample of 23 fundamental mode Cepheids with periods P in the range of $P_0 \pm 1.1$ day, where $P_0 = 7.5721$ day is the period of V440 Per. To this sample we fitted a straight line $\phi_{21}(P) = a(P - P_0) + b$. With this procedure, we find that at the period of V440 Per the expected ϕ_{21} of the fundamental mode Cepheid is 3.628 rad, with the average scatter




Figure 4. Residuals from the mean fundamental mode $P - \phi_{21}$ relation *vs.* pulsation amplitude A_1 . Observational data points taken from Kienzie et al. (1999) and Moskalik et al. (2009). Same symbols as in Fig. 3.

of individual values of $\sigma_0 = 0.026$ rad. This estimate of the intrinsic scatter is conservative, as the nominal ϕ_{21} measurement errors would account for at least half of it. The ϕ_{21} value measured for V440 Per is 2.759 rad, with an error of $\sigma_1 = 0.117$ rad. The probability distribution of the ϕ_{21} offset, Δ , is obtained by convolution of two normal distributions, $N(0, \sigma_0)$ and $N(\Delta, \sigma_1)$. It may be demonstrated, that the result is another normal distribution $N(\Delta, \sqrt{\sigma_0^2 + \sigma_1^2})$, essentially by virtue of the law of error combination. By substituting $\Delta = 3.628 - 2.759 = 0.869$ rad, we find that the observed velocity ϕ_{21} of V440 Per deviates from the fundamental mode sequence by 7.25σ . Thus, from the purely observational evidence we conclude that V440 Per does not pulsate in the fundamental mode. Consequently, it must be an overtone Cepheid.

The argument presented above critically depends on the assumption that velocity ϕ_{21} (at a given period) does not depend on the pulsation amplitude, which for V440 Per is lower than for the fundamental mode Cepheids used as comparison. Such an assumption is well justified by numerical computations (*i.e.* Buchler et al. 1990; Smolec & Moskalik 2008), nevertheless it should be verified with available data. In Fig. 4 we plot residuals from the mean fundamental mode $P - \phi_{21}$ relation *vs.* amplitude A_1 . In this case the mean relation is defined as the parabola fitted to all fundamental mode Cepheids displayed in Fig. 3. For A_1 in the range of 10 – 17 km/s there is no correlation of ϕ_{21} residuals with pulsation amplitude. For $A_1 < 10$ km/s there is a weak evidence of ϕ_{21} increase. As the data are very scarce, we are not convinced that this increase is significant. If it were real, however, it would only strengthen our conclusion that V440 Per strongly deviates from the fundamental mode sequence.

5 MODELING OF V440 PER

Classification of V440 Per as a first overtone pulsator, discussed in the previous Section, is based solely on the morphological properties of Cepheid velocity curves. This mode

Table 3. Convective parameters of pulsation models discussed in this paper. α_s , α_c , α_d , α_p and γ_r are given in the units of standard values (see Smolec & Moskalik 2008). In the last column we give a linear upper limit for first overtone period ($P_{\max}$).

Set	α	α_m	α_s	α_c	α_d	α_p	α_t	γ_r	$P_{\max}$
A	1.5	0.1	1.0	1.0	1.0	0.0	0.00	0.0	8.1 ^d
B	1.5	0.5	1.0	1.0	1.0	0.0	0.00	1.0	8.6 ^d
C	1.5	0.1	1.0	1.0	1.0	1.0	0.01	0.0	13.5 ^d

identification can be strengthened by comparing the velocity curve of V440 Per with those of hydrodynamical Cepheid models. Such a comparison was first performed by Kienzie et al. (1999), who used the unpublished radiative models of Schaller & Buchler (1994). They showed that the theoretical progression of velocity ϕ_{21} supported the overtone classification of V440 Per. However, their conclusion was somewhat weakened by a large error of then available ϕ_{21} of this star.

In the present section, we confirm and extend the results of Kienzie et al. (1999). It is easy to show, that the velocity curve of V440 Per is incompatible with the fundamental mode Cepheid models. Indeed, all published models display velocity ϕ_{21} higher than 3.0 rad *at all periods* and higher than 3.5 rad at the period of V440 Per (*i.e.* Buchler et al. 1990; Moskalik et al. 1992; Smolec & Moskalik 2008). This holds true both for the convective models and for the older radiative models. The computed velocity ϕ_{21} is very robust and shows no sensitivity to the treatment of convection, choice of opacities or to details of the numerical code. Most importantly, it is insensitive to the pulsation amplitude (*cf.* Figs. 8, 11 of Smolec & Moskalik 2008). Clearly, the only chance to match the observed velocity curve of V440 Per is to search for an appropriate overtone model.

With this goal in mind, we computed several sequences of the convective overtone Cepheid models. We show that V440 Per fits theoretical ϕ_{21} progression of the first overtone Cepheids and that its velocity Fourier parameters can be accurately reproduced with hydrodynamical computations.

Modeling of such a long period overtone pulsator is not an easy task. The satisfactory models have to reproduce Fourier parameters and the long period of this variable. The current hydrocodes used for modeling of the radial pulsations adopt a time-dependent convection models (e.g. Stellingwerf 1982, Kuhfuß 1986). These models introduce several dimensionless parameters, that should be adjusted to match the observational constraints. V440 Per with its exceptionally long period, offers an opportunity to impose interesting limits on the free parameters of the convection treatment.

In our computations we use the convective hydrocode of Smolec & Moskalik (2008). The code adopts Kuhfuß (1986) dynamical convection model reformulated for use in stellar pulsation calculations. Kuhfuß model is physically well motivated and self-consistent. It contains 8 scaling parameters, summary of which, together with the detailed description of the model equations is provided by Smolec & Moskalik (2008). The values of parameters used in the present paper are given in Table 3. All our Cepheid models are constructed




Figure 5. Theoretical instability strips for convective parameters of set A.

in the way described in Smolec & Moskalik (2008). In all computations we use Galactic chemical composition ($X = 0.70$, $Z = 0.02$) and OPAL opacities (Iglesias & Rogers 1996) computed with the Grevesse & Noels (1993) mixture. We use the mass-luminosity ($M-L$) relation derived from Schaller et al. (1992) evolutionary tracks ($\log(L/L_\odot) = 3.56 \log(M/M_\odot) + 0.79$).

5.1 Pulsation period of V440 Per

Theoretical instability strips (IS) computed with the linear pulsation code for set A of the convective parameters are presented in Fig. 5. Instability strips for the first overtone and for the fundamental mode are enclosed with the thick and thin lines, respectively. The dotted lines correspond to constant values of the first overtone period, as indicated in the Figure.

First overtone IS does not extend toward an arbitrarily high luminosity. The linear computations yield an upper limit for the first overtone pulsation period, reaching ~ 8.1 day in Fig. 5 (see also Table 3). This limit depends mostly on the adopted convective parameters, but also on the $M-L$ relation and on the metallicity. However, to find a model, that satisfies V440 Per period constraint, it is not enough to assure appropriate linear period limit ($P > 7.5$ day). This is because the maximum overtone period at *full amplitude pulsation* is determined by the nonlinear effects.

Between the linear blue edges of the first overtone and of the fundamental mode and below their crossing point, the first overtone pulsation is the only possibility. As one can see in Fig. 5, such a region is very narrow. A significant part of the overtone IS lays inside the fundamental mode IS. For higher luminosity and overtone periods longer than 6 days, the instability strips of both modes entirely overlap.

In the region, where both modes are linearly unstable,

the final pulsation state is determined by the nonlinear effects (modal selection). Full amplitude pulsation in one of the modes usually suffices to saturate the instability. In principle, a double-mode pulsation is also possible, but it is very unlikely at long overtone periods that are of interest in this paper (no double-mode pulsator with such a long overtone period is known either in the Galaxy, or in the Magellanic Clouds; see Soszyński et al. 2008, Moskalik & Kołaczowski, 2008). Therefore, construction of models matching period of V440 Per requires both linear and nonlinear computations and can be used to constrain parameters of convection model we use.

It is hard to constrain all 8 parameters entering the model. As was shown by Yecko et al. (1998), different combinations of the convective α parameters may yield essentially the same results. Therefore, we decided to freeze four parameters, for which standard values are in use. In all studied sets of the convective parameters (Table 3), mixing-length is set to $\alpha = 1.5$ and for α_s , α_c and α_d we use the standard values (see Smolec & Moskalik 2008 for details). Set A represents the simplest convection model, without turbulent pressure, turbulent flux or radiative losses. These effects are turned on in set B (radiative losses) and in set C (turbulent pressure and turbulent flux). One of the crucial factors of the convection model is eddy-viscous damping, the strength of which is determined by the α_m parameter. The lower the eddy-viscous damping is (lower α_m), the more linearly unstable become the models and, consequently, the higher are their pulsation amplitudes.

The most interesting outcome of the linear computations are the period limits, $P_{\max}$, given in the last column of Table 3. At the linear theory level, all sets of convective parameters listed in the Table can satisfy V440 Per period constraint. This was assured by adjusting eddy-viscous damping parameter, α_m . In case of sets A and C (no radiative losses) low values of α_m are required.

For each parameter set we computed a sequence of nonlinear models, running parallel to the first overtone blue edge, at a constant distance ΔT . In this Section we use $\Delta T = 75$ K. This choice is arbitrary, but we believe, that such models should reproduce most of the observed first overtone variables. The static models were initialized in the first overtone mode (as described in Smolec & Moskalik 2008) and their time evolution was followed, until the final pulsation state was reached. The computed radial velocity curves were then decomposed into Fourier series. In Fig. 6 we plot amplitude A_1 vs. pulsation period for three sequences of models. The most satisfactory results are obtained for set B of the convective parameters. The pulsation amplitudes for this sequence are moderate and they match well the observations (this conclusion will be strengthened in Sect. 5.2). Also, for this sequence we obtain the overtone models with the longest periods. Sequences computed with the parameter sets A and C extend to periods not exceeding 6 days. All more massive models (*i.e.* with longer overtone periods) switched into the fundamental mode pulsation and therefore are not plotted in the Figure. The amplitudes of the models computed with the parameter sets A and C are larger. This is due to the adopted low value of the eddy-viscous damping, needed to obtain high enough limit for the linear overtone period. Any further decrease of α_m does not help. Although $P_{\max}$ increases, yet models switch into the funda-




Figure 6. Fourier amplitude A_1 for three sequences of overtone Cepheid models. Model amplitudes scaled by constant projection factor of 1.4. Except of V440 Per, observational data points are taken from Kienzle et al. (1999) and Moskalik et al. (2009).

mental mode at periods well below of 7.57 days required to match V440 Per. Additionally, at the short periods the amplitudes of the first overtone models become unacceptably high. We conclude, that inclusion of the radiative losses is necessary to reproduce the long overtone period of V440 Per. As we will demonstrate in Sect. 5.2, with the set B of the convective parameters we can also reproduce the Fourier parameters of V440 Per, as well as the overall progression of Fourier parameters for all observed overtone Cepheids.

Our conclusion is consistent with the results of Szabó et al. (2007). For convection with the radiative losses¹, their models extend toward high overtone periods (parameter set A of Szabó et al. 2007). Without radiative losses (set B of Szabó et al. 2007) and with much higher eddy-viscous dissipation than in our set A, their longest overtone periods fall below 3 days. Comparing our models with the overtone Cepheid models of Feuchtinger et al. (2000) we note some inconsistency. Feuchtinger et al. (2000) obtain long overtone periods for all sets of convective parameters considered in their paper. Their results are inconsistent with the results of Szabó et al. (2007) (set A of Feuchtinger et al. 2000 and set B of Szabó et al. 2007 are identical, but their linear results are significantly different) as well as with our results. We trace this discrepancy to different evaluation of the superadiabatic gradient, $Y = \nabla - \nabla_a$, in our and in the Vienna code used by Feuchtinger et al. (2000). In the Vienna code, the radiative pressure contribution is neglected in computation of ∇_a (Feuchtinger 1999). This leads to higher values of ∇_a (≈ 0.4 far from partial ionization regions, see Figs. 3 and 4 of Wuchterl & Feuchtinger 1998) and lower values of the superadiabatic gradient. Consequently, in their models convection is less pronounced and the models are more unstable. In fact, we were able to reproduce their linear results when we neglected the contribution of radiation in the evaluation of ∇_a .

5.2 Fourier parameters of V440 Per

In this Section we focus our attention on set B of the convective parameters. As demonstrated in the previous Section,

¹ Note however, that in Florida-Budapest hydrocode used by Szabó et al. (2007), radiative losses are modeled in a different way (see also Smolec & Moskalik 2008).




Figure 7. Fourier parameters of Cepheid radial velocity curves. Model sequences are computed at constant distances from the first overtone blue edge, as indicated in the Figure. Except of V440 Per, observational data points are taken from Kienzle et al. (1999) and Moskalik et al. (2009). Error bars for most of the amplitudes are smaller than the symbol size and are not plotted.

this parameter set allows to reproduce the long overtone periods and the observed pulsation amplitudes. In Fig. 7 we compare the velocity Fourier parameters of Galactic overtone Cepheids with those of the hydrodynamical models computed with the parameters of set B. Four sequences of models displayed in the plot run at different distances from the first overtone blue edge ($\Delta T = 25$ K, 75 K, 125 K, 175 K). The overall observed progression of Fourier parameters is well reproduced with our models. For ϕ_{21} , we notice that the model sequences are shifted toward somewhat shorter periods. This can be easily explained. The characteristic progression of velocity ϕ_{21} with period is caused by the 2:1 resonance between the first overtone and the fourth overtone (Kienzle et al. 1999; Feuchtinger et al. 2000). The center of this resonance is located at $P = 4.2 - 4.6$ day. The exact location of the resonance in hydrodynamic models depends mostly on the chosen mass-luminosity relation. This relation was not adjusted in our calculations to match the observed resonance progression². As shown in Fig. 5, the resonance center (displayed with the dashed line) crosses the center of the first overtone IS for periods shorter than 4 days, which explains the horizontal shift in Fig. 7. Taking into account this shift, amplitudes, amplitude ratios and Fourier phases agree satisfactorily with observations, although at the short

² Adjustment of $M - L$ relation to satisfy the resonance constraint is not an easy task. Both the slope and the zero point of the adopted $M - L$ relation affect location of the resonance and the shape of the ϕ_{21} progression. In a forthcoming publication (Smolec & Moskalik, in preparation) we address this problem in detail.

periods theoretical ϕ_{21} values are slightly too high. Considering V440 Per, we see the very good agreement with the model sequence closest to the blue edge of the overtone instability strip ($\Delta T = 25$ K). An exact match can be easily obtained for the model sequence located slightly closer to the blue edge. We note, that the progression of the overtone ϕ_{21} at long periods is insensitive to the choice of ΔT . The models predict slightly higher ϕ_{21} than actually observed in V440 Per, but the two values are consistent within the error bar.

6 CONCLUSIONS

Using our new Poznań Spectroscopic Telescope we obtained 158 high precision radial velocity observations of a low amplitude Cepheid V440 Per. We constructed the pulsation velocity curve of V440 Per and we were able to reliably detect its 1st harmonic, with amplitude of only 140 ± 15 m/s. The measured Fourier phase $\phi_{21} = 2.76 \pm 0.12$ rad differs from the values observed in fundamental mode Cepheids of a similar period by 7.25σ . Thus, we demonstrated on purely morphological ground that V440 Per does not pulsate in the fundamental mode. This settles the dispute between Szabó et al. (2007) and Kienzie et al. (1999) and allows to classify V440 Per as an overtone pulsator, the one with the longest period identified so far ($P = 7.57$ day).

Our results demonstrate that with suitable care, our inexpensive instrument featuring MUSICOS Echelle spectrograph and a small robotic telescope can achieve stability and precision surpassed only in the extrasolar planet searches. Note, that we employed neither the iodine cell nor environment control. Yet, our observations prove that the secure mode identification is feasible even for the very low amplitude Galactic Cepheids.

The overtone pulsation of V440 Per has interesting theoretical consequences. To investigate them we employed our convective linear and nonlinear pulsation codes (Smolec & Moskalik 2008). The first overtone linear models are already constrained by the value of the pulsation period at the center of the $2\omega_1 = \omega_4$ resonance (Kienzie et al. 1999; Feuchtinger et al. 2000). The very existence of V440 Per imposes additional constraints. Namely, to linearly excite the first overtone and then to obtain a *nonlinear full-amplitude* overtone pulsation of such a long period, one has to fine tune the dynamical convection model used in the pulsation calculations. Our numerical experiments demonstrate that in the convective energy transport radiative losses must be properly accounted for to maintain consistency with the observations. With this effect taken into account, the nonlinear overtone Cepheid models not only reproduce the exceptionally long period of V440 Per, but they also reproduce neatly all Fourier parameters of its pulsation velocity curve. No such agreement can be achieved with models pulsating in the fundamental mode. These results of the hydrodynamical modeling provide additional support for our empirical classification of V440 Per as a first overtone pulsator.

ACKNOWLEDGMENTS

This publication is based on the first results from Poznań Spectroscopic Telescope, a large instrumental project developed at A. Mickiewicz University in Poznań in the course of many years. Recent work by RB, MF and ASC was supported from the Polish MNiI grant N N203 3020 35. We acknowledge with gratitude permission to use facilities of and help from the Borowiec Geodynamical Observatory of Centre for Space Research (CBK PAN, Borowiec), in general, and encouragement from Prof. Stanisław Schillak, in particular. RS and PM work on the pulsation codes is supported by the MNiSW Grant 1 P03D 011 30. We would like to acknowledge inspiration and support by late Prof. Bohdan Paczynski. Dr. Jacques Baudrand kindly supplied us with the blueprints of MUSICOS.

REFERENCES

- Antonello, E., Poretti, E. & Reduzzi, L., 1990, A&A, 236, 138
- Arellano Ferro, A., 1984, MNRAS, 209, 481
- Baudrand, J. & Bohm, T., 1992, A&A, 259, 711
- Burki, G. & Benz, W., 1982, A&A, 115, 30
- Buchler, J.R., Moskalik, P. & Kovács, G., 1990 ApJ, 351, 617
- Feuchtinger, M., 1999, A&AS, 136, 217
- Feuchtinger, M., Buchler, J.R. & Kolláth, Z., 2000, ApJ, 544, 1056
- Gorynya, N.A., Irmambetova, T.R., Rastorguev, A.S. & Samus, N.N., 1992, Pisma Astron. Zh., 18, 777
- Gorynya, N.A., Samus, N.N., Rastorguev, A.S. & Sachkov, M.E., 1996, Pisma Astron. Zh., 22, 198
- Gorynya, N.A., Samus, N.N., Sachkov, M.E., Rastorguev, A.S., Glushkova, E.V. & Antipin, S.V., 1998, Pisma Astron. Zh., 24, 939
- Grevesse, N. & Noels, A., 1993, in Origin and Evolution of the Elements, Eds. N. Pratz, E. Vangioni-Flam, & M. Casse, Cambridge Univ. Press., p.15
- Iglesias, C.A. & Rogers, F.J., 1996, ApJ, 464, 943
- Kienzie, F., Moskalik, P., Bersier, D. & Pont, F., 1999, A&A, 341, 818
- Kuhfuß, R., 1986, A&A, 160, 116
- Moskalik, P., Buchler, J.R. & Marom, A., 1992, ApJ, 385, 685
- Moskalik, P., Gorynya, N.A. & Samus, N.N., 2009, in preparation
- Moskalik, P. & Kołaczowski, Z., 2008, MNRAS, in press; arXiv:0809.0864
- Ogloza, W., Moskalik, P. & Kanbur, S., 2000, in The Impact of Large-Scale Surveys on Pulsating Star Research, Eds. L. Szabados & D. Kurtz, ASP Conf. Ser. 203, 235
- Schaller, G., Schaerer, D., Meynet, G. & Maeder, A., 1992, A&AS, 96, 269
- Smolec, R. & Moskalik, P., 2008, Acta Astr., 58, 193
- Soszyński, I. et al., 2008, Acta Astr., 58, 153
- Stellingwerf, R.F., 1982, ApJ, 262, 330
- Szabó, R., Buchler, J.R. & Bartee, J., 2007, ApJ, 667, 1150
- Wuchterl, G. & Feuchtinger, M.U., 1998, A&A, 340, 419
- Yecko, P.A., Kolláth, Z. & Buchler, J.R., 1998, A&A, 336, 553

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.

APPENDIX A: FOURIER COEFFICIENTS AND THEIR ERRORS

It seems useful to collect in one place all the relevant formulae for Fourier coefficients of velocity $v(t)$ and their errors σ . Defining Fourier series in the following way

$$\begin{aligned} v(t) - a_0 &= \sum_{n=1}^N A_n \sin(n\omega t + \phi_n) \\ &= \sum_{n=1}^N (a_n \cos n\omega t + b_n \sin n\omega t) \end{aligned} \quad (\text{A1})$$

where N is the order of the fit, we obtain

$$A_n = \sqrt{a_n^2 + b_n^2} \quad \text{and} \quad R_{n1} \equiv \frac{A_n}{A_1} \quad (\text{A2})$$

$$\tan \phi_n = \frac{a_n}{b_n} \quad \text{and} \quad \phi_{n1} \equiv \phi_n - n\phi_1 \quad (\text{A3})$$

From variations of these equations we find

$$\delta A_n = \frac{a_n \delta a_n + b_n \delta b_n}{A_n} \quad (\text{A4})$$

$$\delta R_{n1} = R_{n1} \left(\frac{a_n \delta a_n + b_n \delta b_n}{A_n^2} - \frac{a_1 \delta a_1 + b_1 \delta b_1}{A_1^2} \right) \quad (\text{A5})$$

$$\delta \phi_n = \frac{b_n \delta a_n - a_n \delta b_n}{A_n^2} \quad (\text{A6})$$

$$\delta \phi_{n1} = \frac{b_n \delta a_n - a_n \delta b_n}{A_n^2} - n \frac{b_1 \delta a_1 - a_1 \delta b_1}{A_1^2} \quad (\text{A7})$$

where RHS are scalar products of the vector $(\delta a_n, \delta b_n, \delta a_1, \delta b_1) \equiv \delta F$ and the corresponding derivatives $(\partial/\partial a_n, \partial/\partial b_n, \partial/\partial a_1, \partial/\partial b_1) \equiv L[\cdot]$. For example $\delta \phi_{n1} = L[\phi_{n1}] \cdot \delta F$ and third component of vector $L[\phi_{n1}]$ is $\partial \phi_{n1}/\partial a_1 = -nb_1/A_1^2$. By the law of error propagation, variance of ϕ_{n1} is

$$\sigma_{\phi_{n1}}^2 \equiv \text{Var}[\phi_{n1}] = L[\phi_{n1}] \times \text{Cov}[a_n, b_n, a_1, b_1] \times L[\phi_{n1}]^T \quad (\text{A8})$$

where $\text{Cov}[\cdot]$ denotes the covariance matrix of raw Fourier coefficients (Eq. A1), $\times$ denotes matrix multiplication and superscript T indicates matrix transpose. Analogous expressions hold for other coefficients. The complete linearized transformation matrix M is obtained by substitution of the L vectors corresponding to Eqs. (A4–A7) as its rows. The full covariance matrix of all Fourier coefficients is seldom needed in practice. It may be obtained from Eq. (A8) by substitution of M for L .